

MYCOTAXON

<http://dx.doi.org/10.5248/130.495>

Volume 130, pp. 495–498

April–June 2015

A new species of *Diplococcium* from the Brazilian semi-arid region

DAVI AUGUSTO CARNEIRO DE ALMEIDA¹, TASCIANO DOS SANTOS SANTA IZABEL¹,
LUÍS FERNANDO PASCHOLATI GUSMÃO¹*, & RAFAEL F. CASTAÑEDA-RUÍZ²

¹Universidade Estadual de Feira de Santana, Departamento de Ciências Biológicas,

Av. Transnordestina s/n, Novo Horizonte, 44036-900, Feira de Santana, Bahia, Brazil

²Instituto de Investigaciones Fundamentales en Agricultura Tropical 'Alejandro de Humboldt' (INIFAT),

Académico Titular de la Academia de Ciencias de Cuba,

Calle 1 Esq. 2, Santiago de Las Vegas, C. Habana, Cuba, C.P. 17200

* CORRESPONDENCE TO: lgusmao@uefs.br

ABSTRACT — An undescribed hyphomycete collected during investigations of microfungi on dead plant material in a semi-arid region of northeast Brazil is described and illustrated as *Diplococcium heteroseptatum* sp. nov. The new fungus is distinguished by differentiated brown verruculose conidiophores and oblong to cylindrical 0–12-heteroseptate (but mostly 1-septate) brown conidia.

KEY WORDS— biodiversity, taxonomy, tropical microfungi

Introduction

During a mycological survey of conidial fungi from the semi-arid region in Bahia State, Brazil, a conspicuous fungus was collected that is clearly related to the genus *Diplococcium* Grove, introduced by Grove (1885) with *D. spicatum* as the type species. The genus comprises 29 species characterized by distinct, erect or funiculose, branched, pigmented conidiophores and blastocatenate, unicellular or septate, pigmented conidia produced on polytretic conidiogenous cells (Silva et al. 2014).

Materials & methods

Samples of bark were collected from March 2007 to November 2008 at Serra da Fumaça (10°74'S 40°36'W) and Serra da Jibóia (12°50'S 39°28'W), Bahia state. Methods for specimen collection, isolation, and morphological examination are provided in Almeida et al. (2014). Holotype and additional specimens were deposited in the Herbarium of the State University of Feira de Santana, Bahia, Brazil (HUEFS).




FIG. 1. *Diplococcium heteroseptatum* (holotype, HUEFS 210432): A–B. General aspect. C–D. Conidiophores, conidiogenous cells, and conidia. E–L. Conidia showing the variation in the number of septa.

Taxonomy

Diplococcium heteroseptatum D.A.C. Almeida, T.S. Santa Izabel,

R.F. Castañeda & Gusmão, sp. nov.

FIGS 1, 2

MYCOBANK MB810699

Differs from all other *Diplococcium* spp. by the wide range of conidial septation.

TYPE: Brazil, Bahia State, Pindobaçu, Serra da Fumaça, on bark of an unidentified plant, 20.XI.2008, coll. D.A.C. Almeida (Holotype: HUEFS 210432).

ETYMOLOGY: from the Greek, *hetero-*, referring to the variable numbers of septa + from the Latin, *-septatum*, referring to the septa.

CONIDIOPHORES distinct, single or grouped to profusely fasciculate, erect, unbranched or rarely branched, straight or flexuous, septate, verrucose or verruculose below, smooth toward the apex, brown, $120\text{--}300 \times 4.5\text{--}9\ \mu\text{m}$. CONIDIOGENOUS CELL polytretic, integrated, terminal, smooth, brown. CONIDIA blastocatenate, ellipsoid to oblong or cylindrical, rounded at the ends, 0–12-septate, but mostly 1-septate, constricted at the somewhat thickened septa, smooth, brown, concolorous, $10\text{--}77 \times 6\text{--}10\ \mu\text{m}$.

ADDITIONAL SPECIMENS EXAMINED: BRAZIL, BAHIA, Serra da Jibóia, on bark of an unidentified plant, 27.III.2007, coll. T.S. Santa-Izabel (HUEFS 210428; 210429; 210430; 210430); 9.IV.2007, coll. T.S. Santa-Izabel (HUEFS 210431).

NOTE: *Diplococcium heteroseptatum* also differs from all other *Diplococcium* species by its profusely fasciculate, verrucose or verruculose conidiophores and the random number of conidial septa observed.

Acknowledgments

The authors thank Dr. Prof. Bryce Kendrick and Dr. De-Wei Li for critically reviewing the manuscript. The authors thank the PPBIO Semi-arid (MCTI –CNPq/proc. 554718/2009-0) for financial support. The first and second authors thank the (CAPES) and (CNPq) for scholarships. LFPG thanks CNPq (proc. 305413/2011-2). RFCR is also grateful to the Cuban Ministry of Agriculture and “Programa de Salud Animal y Vegetal”, project P131LH003033, for facilities. We acknowledge the assistance provided by Dr. P.M. Kirk and Drs. V. Robert and A. Decock through the IndexFungorum and MycoBank websites. Dr. Lorelei Norvell’s editorial and Dr. Shaun Pennycook’s nomenclatural reviews are greatly appreciated.

Literature cited

- Almeida DAC, Miller AN, Gusmão LFP. 2014. New species and new combinations of conidial fungi from semi-arid Caatinga biome of Brazil. *Nova Hedwigia* 98: 431–447. <http://dx.doi.org/10.1127/0029-5035/2013/0162>
- Grove WB. 1885. New or noteworthy fungi. II. *Journal of Botany British and Foreign*. 23: 155–168.
- Silva SS, Cruz ACR, Gusmão LFP, Castañeda-Ruiz RF. 2014. *Diplococcium variegatum* a new conidial fungi from semi-arid Caatinga biome of Brazil. *Mycotaxon* 127: 59–62. <http://dx.doi.org/10.5248/127.59>




FIG. 2. *Diplococcium heteroseptatum* (holotype, HUEFS 210432):
A: Conidiophores, conidiogenous cells, and conidia. B: Conidiogenous cell and conidia.